

On the displacement rigidity of Levi flat hypersurfaces - The case of boundaries of disc bundles over compact Riemann surfaces ^{*†}

Klas Diederich Takeo Ohsawa

April 15, 2005

Abstract

Non-existence theorems for Levi flat hypersurfaces have found great interest in the literature. The question next to this that has to be asked is, when existing Levi flat hypersurfaces are at least rigid under deformations. Here, the case of boundaries of disc bundles over certain compact Riemann surfaces is considered.

Introduction

In several complex variables, Levi flat hypersurfaces arose as counterexamples in generalized function theory on complex manifolds (cf. [12, 13, 1, 3]). Nowadays, they are considered as objects of independent interest, since they also arise as typical examples of minimal closed subsets consisting of leaves of complex analytic foliations (cf. [2]). However, very few Levi flat hypersurfaces have yet been analyzed. The most remarkable results are nonexistence theorems (cf. [17],[18],[9]). Recently an attempt has begun to classify them (cf. [11], [14]). Under these circumstances we would like to continue the study of disc bundles over compact Kähler manifolds from [5], where we proved as main result that any holomorphic disc bundle D over a compact Kähler manifold M is weakly 1-complete (i.e. it admits a $\mathcal{C}^\infty$ -plurisubharmonic exhaustion function).

What we want to pursue further, is a rigidity property of ∂D when D is identified with a domain in the associated $\mathbb{P}^1$ -bundle, say $P \rightarrow M$. For that we shall restrict ourselves here, as a first step, to the case where M is a compact Riemann surface of genus $g > 1$, since we can exploit some (deep) results on Riemann surfaces.

^{*}MSC 2000: 32G07, 32G08, 32V15, 32L05, 30F35, 14H30

[†]Keywords: Levi flats, rigidity, disc bundles, compact Riemann surfaces

First, employing Schoen-Yau's diffeomorphism theorem for harmonic maps, we refine the previous result as follows.

Proposition 0.1 (Consequence of Proposition 1.6) *Let C be a compact Riemann surface of genus $g > 1$ and let $D \rightarrow C$ be a holomorphic disc bundle associated to a homomorphism ϱ from the fundamental group of C into the automorphism group of the unit disc $\mathbb{D}$. If the image Γ of ϱ is a Fuchsian group such that $\mathbb{D}/\Gamma$ is homeomorphic to C , then D is Takeuchi 1-convex in the associated $\mathbb{P}^1$ -bundle P . (For the definitions see §2)*

Based on this observation, we can conclude the following rigidity result.

Theorem 0.2 *∂D is rigid in P if either Γ is an abelian group or a Fuchsian group such that $\mathbb{D}/\Gamma$ is biholomorphic to C or to its conjugate $\overline{C}$.*

(For the definition of rigidity, see §2.) For the proof of the theorem we need a Hartogs type extension theorem of Ivashkovitch [10] and a basic fact on projective structures described by R. C. Gunning in [7, 8].

The condition on Γ does not seem to be really essential for the rigidity of ∂D . One might even suspect that the rigidity holds true for any disc bundle over any compact complex manifold. Although there are few methods available to study the question in such generality, the authors believe that the present work gives some insight towards that direction.

Acknowledgement: The second-named author would like to express his gratitude to the university of Wuppertal for the hospitality during the preparation of this work. The authors are very grateful to the referee for communicating to them equality (6) which they were not aware of.

1 Disc bundles over compact Kähler manifolds - review and refinement

Let M be a compact complex manifold of dimension n and let $D \rightarrow M$ be a holomorphic fiber bundle the fibers of which are biholomorphic to the unit disc $\mathbb{D} := \{\zeta \in \mathbb{C} : |\zeta| < 1\}$. Recall that the group $\text{Aut } \mathbb{D}$ of biholomorphic automorphisms of $\mathbb{D}$ consists of the maps

$$\zeta \mapsto e^{i\theta} \frac{\zeta - a}{\bar{a}\zeta - 1} \quad (\theta \in \mathbb{R}, a \in \mathbb{D})$$

so that transition maps of the bundle D are locally constant. Accordingly, the pull-back of D to the universal covering $\tilde{M} \rightarrow M$ is the trivial disc bundle $\tilde{M} \times \mathbb{D}$, and there exists a homeomorphism

$$\varrho : \pi_1(M, x_0) \rightarrow \text{Aut } \mathbb{D} \quad (x_0 \in M) \tag{1}$$

uniquely determined up to the inner automorphism operation of $\text{Aut } \mathbb{D}$, such that

$$D \simeq \tilde{M} \times \mathbb{D} / \sim \tag{2}$$

where $(x, \zeta) \sim (y, \zeta') : \Leftrightarrow y = \sigma(x)$ and $\zeta' = \varrho(\sigma)\zeta$ for some $\sigma \in \pi_1(M, x_0)$. Here the action of σ is defined as the covering transformation. We shall denote D by D_ϱ when we want to refer to ϱ . In particular, D is uniquely determined by ϱ .

Let $P \rightarrow M$ be the $\mathbb{P}^1$ -bundle associated to the composite of ϱ and the inclusion homomorphism $\text{Aut } \mathbb{D} \hookrightarrow \text{Aut } \mathbb{P}^1$. D will be naturally identified with a domain in P .

Proposition 1.1 *For any compact Kähler manifold M and for any disc bundle D over M , one of the following four cases occurs:*

1. D admits a unique locally nonconstant pluriharmonic section.
2. D admits a locally constant section.
3. $\partial D \rightarrow M$, the associated circle bundle, admits a unique locally constant section.
4. $\partial D \rightarrow M$ admits precisely two locally constant sections.

Proof: It is clear that 2) occurs if and only if D is the tubular neighborhood of the zero section of a topologically trivial line bundle over M . 3) (resp. 4)) occurs if and only if the transition maps have one (resp. two) common fixed points on the boundary of $\mathbb{D}$, in which case there are no (resp. infinitely many) pluriharmonic sections of D . By [5] the rest is contained in the (possibly empty) case 1). $\square$

Corollary 1.2 (cf. [5]) *Any holomorphic disc bundle over a compact Kähler manifold is weakly 1-complete.*

Definition 1.3 *A $\mathcal{C}^2$ real valued function φ on a complex manifold X of dimension n is said to be q -convex at a point $x \in X$ if the Levi form of φ has at least $n - q + 1$ positive eigenvalues at x .*

In the cases 1), 3) and 4) of Proposition 1.1 for the disc bundles D over a compact Kähler manifold of dimension n , it turns out that D admits an exhaustion function of class $\mathcal{C}^\infty$ which is n -convex outside a compact subset of D . This can easily be seen from the proof of the above Corollary which we have given in [5].

Instead of this general fact, we shall prove a refined variant which we shall need later.

Definition 1.4 *A relatively compact domain Ω with $\mathcal{C}^2$ -smooth boundary in a complex manifold X is said to be Takeuchi q -convex if Ω admits a defining function r of class $\mathcal{C}^2$ such that, with respect to some Hermitian metric on X , at least $n - q + 1$ eigenvalues of the Levi form of $-\log(-r)$ are greater than 1 outside a compact subset of Ω .*

Remark 1.5 *A. Takeuchi [19] was the first to verify that the q -convexity in the above sense holds for $q = 1$, if $X = \mathbb{P}^n$ and Ω is a proper locally pseudoconvex domain.*

Proposition 1.6 *Let C be a compact Riemann surface of genus ≥ 2 , and let $D \rightarrow C$ be a disc bundle with a harmonic section $s : C \rightarrow D$. Suppose that the set of critical points of s is finite. Then D admits a defining function r in the associated $\mathbb{P}^1$ -bundle P such that $\partial\bar{\partial}(-\log(-r))$ dominates the ambient metric near ∂D . In other words, D is Takeuchi 1-convex in P .*

Proof: In the above situation we define a function φ on D by putting

$$\varphi(x, \zeta) := -\log \left(1 - \left| \frac{\zeta - h(x)}{\overline{h(x)}\zeta - 1} \right|^2 \right)$$

in terms of the coordinates ζ on the fibers.

Clearly, the value φ does not depend on the choice of the coordinates ζ , and φ is a real analytic exhaustion function of D .

Let x_0 be any point of C and let ζ be a coordinate on the fiber D_{x_0} satisfying $h(x_0) = 0$. Then a simple calculation gives with respect to a local coordinate z around x_0 :

$$\varphi_{z\bar{z}} = |h_z|^2 + |h_{\bar{z}}|^2 - 2\operatorname{Re}(\bar{\zeta}^2 h_z h_{\bar{z}}) \quad (3)$$

$$\varphi_{\zeta\bar{\zeta}} = (1 - |\zeta|^2)^{-2} \quad (4)$$

$$\varphi_{z\bar{\zeta}} = -h_z \quad (5)$$

at (x_0, ζ) . Hence

$$\varphi_{z\bar{z}}\varphi_{\zeta\bar{\zeta}} - \left| \varphi_{z\bar{\zeta}} \right|^2 = (1 - |\zeta|^2)^{-2} \left[|\zeta|^2 (1 - |\zeta|^2) |h_z|^2 + |\bar{\zeta} h_z - \overline{h_{\bar{z}}}|^2 \right] \quad (6)$$

holds true.

Let $\{x_1, \dots, x_m\}$ be the set of critical points of s , and let ζ_i be fiber coordinates of D over neighborhoods U_i of x_i .

Let ϱ_i be a C^∞ nonnegative functions on C whose supports are contained in U_i , such that ϱ_i is identically 1 near x_i .

We put, for $\varepsilon > 0$,

$$\Phi = \varphi + \varepsilon \sum_{i=1}^m \left\{ -\varrho_i \log(1 - |\zeta_i|^2) + (1 - \varrho_i)\varphi \right\}$$

Then (4) and (6) imply that there exists a Hermitian metric g_P on the associated bundle $P \rightarrow C$ such that $\partial\bar{\partial}\Phi > g_P|D$ holds outside a compact subset of D , if ε is chosen to be sufficiently small.

Remark 1.7 Notice that for a disc bundle $D \rightarrow C$ associated to the homomorphism $\varrho : \pi_1(C, x_0) \rightarrow \operatorname{Aut} \mathbb{D}$, the exterior of D in the associated $\mathbb{P}^1$ -bundle $P \rightarrow C$ is equivalent to the disc bundle associated to the homomorphism $\bar{\varrho}$ defined by

$$\bar{\varrho}(\sigma)(z) := \overline{\varrho(\sigma)(\bar{z})}$$

Therefore, if $(x, h(x))$ is a locally quasiconformal section of D , $(x, \overline{h(x)})$ is a locally ant quasiconformal section of $D_{\bar{\varrho}}$.

In order to deduce the Proposition of the introduction from Proposition 1.6, we note that a $\mathcal{C}^\infty$ -section of $D_\varrho \rightarrow C$ is naturally identified with a $\mathcal{C}^\infty$ -map from C to $\mathbb{D}/\text{Im } \varrho$ which is homotopic to a diffeomorphism. Therefore, the harmonic section of $D_\varrho \rightarrow C$, which exists according to the theorem of Eells and Sampson (cf. [6]), is either quasiconformal or ant quasiconformal as a map to $\mathbb{D}/\text{Im } \varrho$, since it is a diffeomorphism in virtue of a theorem of Schoen-Yau [15]. Hence, the required Takeuchi 1-convexity follows from Proposition 1.6.

2 Stability of q -convexity

Before starting to discuss the rigidity property of Levi-flat hypersurfaces, we shall prove the stability of Takeuchi q -convexity for domains with Levi-flat boundaries.

Recall that a $\mathcal{C}^2$ -smooth real hypersurface S in a complex manifold X is said to be Levi-flat if S locally admits a defining function the Levi form of which, restricted to the holomorphic tangent space of S , is identically 0. A real hypersurface of class $\mathcal{C}^\omega$ then is defined by pluriharmonic functions if and only if it is Levi-flat.

Proposition 2.1 *Let $\Omega \subset\subset X$ be a Takeuchi q -convex domain with a Levi-flat $\mathcal{C}^\omega$ hypersurface as boundary. Then any $\mathcal{C}^2$ small perturbation of Ω as a domain with $\mathcal{C}^\omega$ Levi-flat boundary also is Takeuchi q -convex. In other words, there exists a tubular neighborhood U and a $\mathcal{C}^\omega$ diffeomorphism φ between U and the normal bundle of $\partial\Omega$ which identifies $\partial\Omega$ with the zero section of the bundle, such that a $\mathcal{C}^\omega$ domain $\Omega' \subset\subset X$ is Takeuchi q -convex if $\partial\Omega'$ is Levi flat and sufficiently small as a $\mathcal{C}^2$ section of the bundle.*

Proof: Let r be a $\mathcal{C}^2$ defining function of $\partial\Omega$ such that $\partial\bar{\partial}(-\log(-r))$ has at least $n-q+1$ eigenvalues > 1 near $\partial\Omega$ with respect to a Hermitian metric on X . Since $\partial\Omega$ is $\mathcal{C}^\omega$ and Levi flat, r is locally the product of a pluriharmonic defining function, say r_α , and a positive $\mathcal{C}^2$ function, say u_α . If Ω' is a domain with $\mathcal{C}^\omega$ Levi flat boundary such that $\partial\Omega'$ is sufficiently close to $\partial\Omega$ in the $\mathcal{C}^\omega$ topology, then one can choose locally pluriharmonic defining functions of $\partial\Omega'$ which are close to r_α even in the $\mathcal{C}^\omega$ topology. Therefore, Ω' admits a defining function r' whose pluriharmonic and positive factors are close to the corresponding factors r_α and u_α in the $\mathcal{C}^2$ topology, respectively.

Combining this observation with

$$\begin{aligned} \partial\bar{\partial}(-\log(-r)) &= \partial\bar{\partial}(-\log(-r_\alpha u_\alpha)) \\ &= \partial\bar{\partial}(-\log(-r_\alpha)) + \partial\bar{\partial}(-\log u_\alpha) \\ &= \frac{\partial r_\alpha \bar{\partial} r_\alpha}{r_\alpha^2} + \partial\bar{\partial}(-\log u_\alpha) \end{aligned}$$

one can immediately see the validity of the conclusion because of the continuity of the eigenvalues of $\partial\bar{\partial}(-\log u_\alpha)$. $\square$

Definition 2.2 *In what follows we say that $\partial\Omega'$ is a displacement of $\partial\Omega$ if $\partial\Omega'$ is identifiable with a section of the normal bundle of $\partial\Omega$ by means of some diffeomorphism φ as*

in Prop. 2.1. "Sufficiently small" always refers to the $\mathcal{C}^2$ -norm. $\partial\Omega$ is said to be rigid if, for any fixed choice of φ , any sufficiently small displacement of $\partial\Omega$ is isomorphic to $\partial\Omega$ as a CR manifold.

Proposition 2.3 *Let $D \rightarrow C$ be a disc bundle as in Proposition 1.6. Then any small $\mathcal{C}^\omega$ Levi flat displacement of ∂D in the associated $\mathbb{P}^1$ bundle $P \rightarrow C$ bounds a Takeuchi 1-convex domain.*

3 Proof of the rigidity theorem

First we will prove case 1) of the Theorem.

Let C be a compact Riemann surface of genus $g \geq 2$ and let $D_\varrho \rightarrow C$ be a disc bundle such that the image Γ of ϱ satisfies $\mathbb{D}/\Gamma \simeq C$ or $\mathbb{D}/\Gamma \simeq \tilde{C}$. By remark 1.7 it suffices to prove the theorem in the case $\mathbb{D}/\Gamma \simeq C$, which we shall now consider.

Since D_ϱ is according to the assumption biholomorphic to the quotient of $\mathbb{D} \times \mathbb{D}$ modulo the diagonal action of Γ by $(z, z) \mapsto (\gamma(z), \gamma(z))$ for $\gamma \in \Gamma$, D_ϱ admits a holomorphic section $s : C \rightarrow D_\varrho$ corresponding to the diagonal $\Delta = \{(z, z) : z \in \mathbb{D}\}$.

Suppose now there exists a sequence of real analytic Levi flat hypersurfaces S_k , $k = 1, \dots$, converging to ∂D_ϱ in the $\mathcal{C}^2$ sense. Then by Proposition 2.3, S_k ($k \gg 1$) separates the associated $\mathbb{P}^1$ bundle P_ϱ into two Takeuchi 1-convex domains, say D_k^+ and D_k^- . We normalize notation such that $D_k^+ \supset s(C)$.

We note that the domain D_k^- does not contain any compact complex curve. In fact, if there were such a curve $C_k \subset D_k^-$, C_k would define a multivalued holomorphic section of the affine line bundle $P_\varrho \setminus s(C)$. By averaging C_k produces a holomorphic section, so that the bundle becomes a line bundle. On the other hand, it is known that ϱ lifts to a $\mathrm{GL}(2, \mathbb{C})$ representation, say $\tilde{\varrho}$, and the rank two vector bundle V associated to $\tilde{\varrho}$ admits a flat connection (cf. [7]). However, s and the zero section of $P_\varrho \setminus s(C)$ lift to line subbundles L_∞ and L_0 of V , so that V is holomorphically equivalent to $L_0 \oplus L_\infty$. This means that, by Weil's criterion on the existence of flat connections (cf. [8]), L_0 and L_∞ both admit flat connections, which is an absurdity because $|\deg L_0| = |\deg L_\infty| = 2g - 2 \neq 0$. Hence D_k^- is Stein.

On the other hand, since S_k is $\mathcal{C}^\omega$ Levi flat, there exists a neighborhood $U_k \supset S_k$ and a complex analytic foliation $\mathcal{F}_k$ on U_k of codimension 1 which extends the foliation on S_k defined by the holomorphic tangent bundle of S_k . We note that $\mathcal{F}_k$ is naturally identified with a holomorphic map from U_k to the projectivization of the tangent bundle of P . Hence, in virtue of the extension theorem of Ivashkovitch (cf. [10]), D^- admits a holomorphic foliation, possibly with finitely many singularities, which extends $\mathcal{F}_k$. Similarly, since D_k^+ does not contain any compact complex curves other than $s(C)$ ($s(C)$ would not be exceptional otherwise), $\mathcal{F}_k$ extends to a holomorphic foliation on $D_k^+ \setminus s(C)$, possibly with finitely many singularities. But since $s(C)$ is exceptional, the foliation further extends, in virtue of the Remmert-Stein continuation theorem for complex analytic subsets (cf. [16]), to D_k^+ , possibly with finitely many singularities. Since these foliations converge to a foliation $\mathcal{F}$ of P_ϱ consisting of locally flat sections, they have for sufficiently large

k no singularities. (Any small perturbation of a holomorphic section as a meromorphic section is holomorphic.)

Thus we obtain a family of holomorphic foliations, say $\tilde{\mathcal{F}}_k$ on P_ϱ , for $k \gg 1$, extending $\mathcal{F}_k$ and converging to $\mathcal{F}$. Note that $\mathcal{F}_k$ defines a flat structure on P_ϱ . Let ϱ_k be the corresponding representation of $\pi_1(C, x_0)$ into $\mathrm{PSL}(2, \mathbb{C})$. Then $\partial D_k^+ \cap P_{x_0}$ must be a circle in the Riemann sphere P_{x_0} because it is a closed simple closed curve which contains an orbit of a point, consisting of infinitely many points, through the action of γ^l ($l = 1, 2, \dots$) for some $\gamma \in \mathrm{PSL}(2, \mathbb{C})$. Therefore, $D_k^\pm$ are biholomorphically equivalent to some disc bundles over C .

This means, that the ϱ_k are equivalent to $\mathrm{Aut}(\mathbb{D})$ -representations of $\pi_1(C, x_0)$, say ϱ'_k . Then ϱ'_k must be $\mathrm{Aut}(\mathbb{D})$ -equivalent to ϱ , because $D_{\varrho'_k}$ and D_ϱ both contain $s(C)$ as a holomorphic section. Therefore, there exist bundle equivalences $\varphi_k : D_\varrho \rightarrow D_{\varrho'_k}$ which extend, by inversion, to bundle automorphisms $\tilde{\varphi}_k$ of $P_\varrho (= P_{\varrho_k} = P_{\varrho'_k})$.

Now we are going to prove case 2) of the Theorem. Since Γ now is abelian, the transition maps of D are either all elliptic, all parabolic or all hyperbolic (if they are not the identity).

Suppose first that either D is trivial or the transition maps are all elliptic, and let $\{\zeta_\alpha\}$ be a system of fiber coordinates of D subordinate to an open covering $\{U_\alpha\}$ of C such that $\zeta_\alpha = e^{i\theta_{\alpha\beta}}$ for $\theta_{\alpha\beta} \in \mathbb{R}$ over $U_\alpha \cap U_\beta$.

Let $\varepsilon > 0$ and let S be any $\mathcal{C}^\omega$ Levi flat displacement of ∂D in the domain $\{1 - \varepsilon < |\zeta_\alpha| < 1 + \varepsilon\}$. Let Ω_+ and Ω_- be the components of $P \setminus S$, such that $\Omega_+ \subset \{|\zeta_\alpha| < \infty\}$. We define a continuous function δ on $\{|\zeta_\alpha| < \infty\}$ by letting

$$\delta(x, \zeta_\alpha) = \inf\{|\zeta_\alpha - \zeta'_\alpha| : (x, \zeta'_\alpha) \in S\} \quad (7)$$

Then $-\log \delta + |\zeta_\alpha|^2$ is plurisubharmonic on Ω_+ and real analytic near S . Since Ω_+ is clearly not 1-convex, $-\log \delta$ must depend only on the fiber coordinate (cf. [4]). Hence S is CR-equivalent to ∂D by the fiberwise retraction along the radial directions with respect to ζ_α .

Next, suppose that the fiber maps are all parabolic, and let $\sigma : C \rightarrow \partial D$ be the locally constant section consisting of the common fixed points of the transition maps. In view of the classification in Proposition 1.1 and the extension argument for D_ϱ with Fuchsian representation ϱ , it suffices to show that any small $\mathcal{C}^\omega$ Levi flat displacements of ∂D are the boundaries of 1-convex domains from both sides. In fact, if $D' \rightarrow C$ is another disc bundle such that the associated $\mathbb{P}^1$ bundle $P' \rightarrow C$ is biholomorphic to P , P and P' are equivalent $\mathbb{P}^1$ bundles because the genus of C is not zero. Hence, the transition maps of D' are all parabolic, too. Hence $\partial D'$ must contain $\sigma(C)$, so that there exists a biholomorphism between D and D' given by the fiberwise translations.

Let Ω_+ and Ω_- be the connected components of $P \setminus S$ which we want to prove to be 1-convex. If $S \supset \sigma(C)$, there is nothing left to prove because $P \setminus \sigma(C)$ is already Stein. So let us suppose $S \not\supset \sigma(C)$. Then $S \cap \sigma(C)$ is either empty or the union of finitely many irreducible real analytic curves, say $\gamma_1, \dots, \gamma_N$, because of the real analyticity of S . If $S \cap \sigma(C) = \emptyset$, $\Omega_+ \supset \sigma(C)$ or $\Omega_- \supset \sigma(C)$. In any case, since $\Omega_\pm \setminus \sigma(C)$ are Stein, there would arise a 2-dimensional Stein manifold with a disconnected boundary which is

absurd.

Hence $S \cap \sigma(C) \neq \emptyset$. Let π be the bundle projection $P \rightarrow C$. Then $\pi(\gamma_k)$ are all real analytic curves, so that $C \setminus \bigcup \pi(\gamma_k)$ carries a bounded strictly subharmonic function say ψ .

On the other hand, let ζ_α be the fiber coordinates of $P \setminus \sigma(C)$ whose transition relations are $\zeta_\alpha = \zeta_\beta + \xi_{\alpha\beta}$, $\xi_{\alpha\beta} \in \mathbb{C}$. Hence we define a function δ on $\Omega_\pm \setminus \sigma(C)$ by $\delta(x, \zeta_\alpha) := \inf\{|\zeta_\alpha - \zeta'_\alpha| : (x, \zeta'_\alpha) \in S\}$ and obtain a plurisubharmonic function φ on $\Omega_\pm$ by extending $-\log \delta$ to $\sigma(C) \setminus S$ as $-\infty$.

Let $d(p)$ be the distance from p to S with respect to some real analytic metric g on P , let $\chi : P \rightarrow [0, 1]$ be a $\mathcal{C}^\infty$ function such that $\chi \equiv 1$ on a neighborhood of $S \cap \sigma(C)$ and that there exists a strictly plurisubharmonic function, say η , on a neighborhood of $\text{supp } \chi$. Then we put

$$\Phi := \max\{\lambda(-\log \delta + \psi), -\log d + \chi\eta\} \quad (8)$$

for a $\mathcal{C}^\infty$ convex increasing function λ with $\inf \lambda = 0$.

Clearly, if S is a sufficiently small displacement of ∂D , Φ is an exhaustion function of $\Omega_\pm$ and satisfies

$$\partial\bar{\partial}\Phi > cg \quad (9)$$

near S for some positive constant c , in the distribution sense. Therefore $\Omega_\pm$ are 1-convex.

Finally, suppose that the transition maps are hyperbolic. Then $P \rightarrow C$ admits two holomorphic curves C_1 and C_2 , lying in ∂D , which consist of the common fixed points of the transition maps. Then the $\mathbb{C}^*$ bundle $P \setminus (C_1 \cup C_2)$ admits a system of fiber coordinates $\{\zeta_\alpha\}$ with transition relations $\zeta_\alpha = e^{i\theta_{\alpha\beta}} \zeta_\beta$ ($\theta_{\alpha\beta} \in \mathbb{R}$). Hence, for any sufficiently small $\mathcal{C}^\omega$ Levi flat displacement S of ∂D , the components of $P \setminus S$ are 1-convex, similar as in the parabolic case. The rest also is similar to that case. Q.E.D.

References

1. Barrett, D.: *Biholomorphic domains with inequivalent boundaries*, Inventiones Math. **85** (1986), 373–377.
2. Cerveau, D.: *Minimaux des feuilletages algébriques de $\mathbb{CP}^n$* , Ann. Inst. Fourier **43** (1993), 1535–1543.
3. Diederich, K., Fornæss, J. E.: *A smooth pseudoconvex domain without pseudoconvex exhaustion*, manuscripta math. **39** (1982), 119–123.
4. Diederich, K., Ohsawa, T.: *A Levi problem on two-dimensional complex manifolds*, Math. Ann. **261** (1982), 255–261.
5. Diederich, K., Ohsawa, T.: *Harmonic mappings and disc bundles over compact Kähler manifolds*, Publ. RIMS Kyoto Univ. **21** (1985), 819–833.

6. Eells, J., Sampson, J. H.: *Harmonic mappings of Riemannian manifolds*, Amer. J. Math. **86** (1964), 109–160.
7. Gunning, R.: *Lectures on Riemann surfaces*, Mathematical Notes, vol. No. 2, Princeton University Press, Princeton, N. J., 1966.
8. Gunning, R.: *Lectures on vector bundles over Riemann surfaces*, Mathematical Notes, vol. No. 6, Princeton University Press, Princeton, N. J., 1967.
9. Iordan, A.: *On the non-existence of smooth Levi-flat hypersurfaces in $\mathbb{CP}^n$* , Proceedings of the Memorial Conference of Kioshi Oka's Centennial Birthday on Complex Analysis in Several Variables, Kyoto, Nara.
10. Ivashkovitch, S.: *The Hartogs-type extension theorem for meromorphic maps into compact Kähler manifolds*, Invent. math. **109** (1992), 47–54.
11. Matsumoto, K., Ohsawa, T.: *On the real analytic Levi flat hypersurfaces in complex tori of dimension two*, Ann. Inst. Fourier **52** (2002), 1525–1532.
12. Narasimhan, R.: *The Levi problem in the theory of functions of several complex variables*, Proc. Internat. Congr. Math. (Stockholm 1962) (Uppsala), Almqvist and Wiksells, pp. 385–388.
13. Ohsawa, T.: *A Stein domain with smooth boundary which has a product structure*, Publ. Res. Inst. Math. Sci. **18** (1982), 1185–1186.
14. Ohsawa, T.: *Levi-flats in complex tori*, In preparation, 2004.
15. Schoen, R., Yau, S. T.: *On univalent harmonic maps between surfaces*, Invent. math. **44** (1978), 265–278.
16. Siu, Y.-T.: *Techniques of Extension of Analytic Objects*, Lecture Notes in Pure and Applied Math., vol. 8, Marcel Dekker, New York, 1974.
17. Siu, Y. T.: *Nonexistence of smooth Levi-flat hypersurfaces in complex projective spaces of dimension ≥ 3* , Ann. Math. **151** (2000), 1217–1243.
18. Siu, Y.-T.: *$\bar{\partial}$ -regularity for weakly pseudoconvex domains in hermitian symmetric spaces with respect to invariant metrics*, Ann. Math. **156** (2002), 595–621.
19. Takeuchi, A.: *Domaines pseudoconvexes infinis et la métrique riemannienne dans un espace projectif*, J. Math. Soc. Japan **16** (1964), 159–181.

Klas Diederich
 Mathematik
 Universität Wuppertal
 Gausstr. 20
 D-42097 Wuppertal
 Germany

Takeo Ohsawa
 Graduate School of Mathematics
 Nagoya University
 Chikusa-ku, Furocho
 Nagoya 464-01
 JAPAN